

A Lifting Theorem for Locally Convex Subspaces of L_0

R. G. Faber

Abstract. We prove that for every closed locally convex subspace E of L_0 and for any continuous linear operator T from L_0 to L_0/E there is a continuous linear operator S from L_0 to L_0 such that $T = QS$ where Q is the quotient map from L_0 to L_0/E .

§0. Introduction.

Let E be a subspace of $L_0 = L_0[0, 1]$, the space of all measurable functions from $[0, 1]$ to $\mathbb{R}$. Let T be an operator from L_0 to L_0/E . What conditions on E ensure that we can find an operator S that makes the following diagram commute?

$$\begin{array}{ccc} & & L_0 \\ & \nearrow S & \downarrow Q \\ L_0 & \xrightarrow{T} & L_0/E \end{array}$$

A. Pełczyński was the first to ask if locally convex subspaces E have this property. If E is locally bounded then we can find such an operator (Kalton - Peck [2]). Peck - Starbird [6] showed that this is also true when E is isomorphic to ω , the space of all real sequences. The goal of this paper is to show that if E is locally convex then we can complete the previous diagram.

We will state some notation. We will let μ represent the standard Lebesgue measure. We also define the map $f \mapsto \|f\|_0$ ($L_0 \rightarrow \mathbb{R}$) as

$$\|f\|_0 = \int_0^1 \frac{|f(x)|}{1 + |f(x)|} dx.$$

This map is an F-norm on L_0 , that is,

- | | | |
|-------|--|------------------------------------|
| (i) | $\ f\ _0 > 0$ | $f \neq 0,$ |
| (ii) | $\ \alpha f\ _0 \leq \ f\ _0$ | $ \alpha \leq 1$ and $f \in L_0,$ |
| (iii) | $\lim_{\alpha \rightarrow 0} \ \alpha f\ _0 = 0$ | $f \in L_0,$ |
| (iv) | $\ f + g\ _0 \leq \ f\ _0 + \ g\ _0$ | $f, g \in L_0.$ |

The map also induces a metric on L_0 . The topology induced by the L_0 metric is just the topology of convergence in measure. For $f \in L_0$ we define $\sigma : L_0 \rightarrow [0, 1]$ by

$$\sigma(f) = \sup_{n \in \mathbb{N}} \|nf\|_0 = \lim_{n \rightarrow \infty} \|nf\|_0.$$

1991 *Mathematics Subject Classification*: Primary 46E30, 46A22.

This paper is part of the author's Ph.D. thesis at the University of Illinois, written under the supervision of N.T. Peck. The research was supported in part by a Beckman grant from the Research Board of the University of Illinois.

By the dominated convergence theorem we can see that $\sigma(f) = \mu(\text{supp } f)$, where $\text{supp } f = \{x : |f(x)| > 0\}$. The F-norm on the quotient space L_0/E is defined in the usual way

$$\|\gamma\|_{L_0/E} = \inf_{f \in \gamma} \|f\|_0 \quad \text{for all } \gamma \in L_0/E.$$

For a subset A of $[0, 1]$ we will let $L_0(A)$ mean the subspace of L_0 consisting of all functions supported on A . We define

$$\|f\|_{L_0(A)} = \|f \cdot \chi_A\|_0,$$

where χ_A is the characteristic function of A .

§1. Preliminary Lemmas.

We have a lifting theorem for locally bounded subspaces (See Theorem 3.6 of [2].) and we will see that locally convex subspaces are in some sense almost locally bounded. The lemmas that follow show us that the ‘unbounded part’ of a locally convex subspace is arbitrarily small. Lemma 1.2 is at the heart of this argument. However, we first need a lemma from Paley and Zygmund [5].

Lemma 1.1. *Let $\alpha > \beta \geq 0$. If $f \in L_0[0, 1]$ such that $\int_0^1 f \geq \alpha$ and $\|f\|_2 = 1$ then*

$$\mu(t : f(t) \geq \beta) \geq (\alpha - \beta)^2.$$

Proof.

$$\begin{aligned} \alpha &\leq \int_0^1 f = \int_{\{f \geq \beta\}} f + \int_{\{f < \beta\}} f \\ &\leq \int_0^1 f \cdot I_{\{f \geq \beta\}} + \beta \\ &\leq \|f\|_2 \cdot \|I_{\{f \geq \beta\}}\|_2 + \beta && \text{(Schwarz Inequality)} \\ &= \sqrt{\mu(t : f(t) \geq \beta)} + \beta. \end{aligned}$$

So $\mu(t : f(t) \geq \beta) \geq (\alpha - \beta)^2$. □

Notice that Rademacher functions do not appear in the statement of the next lemma but they play a key role in the proof. Recall that all the Rademacher functions act on $[0, 1]$ and have values in the two point set $\{-1, 1\}$. The first Rademacher function, r_1 , is 1 everywhere. The second, r_2 , is 1 on $[0, \frac{1}{2})$ and -1 on $[\frac{1}{2}, 1]$; r_3 is 1 on $[0, \frac{1}{4})$ and $[\frac{1}{2}, \frac{3}{4})$ but -1 on $[\frac{1}{4}, \frac{1}{2})$ and $[\frac{3}{4}, 1]$; and so on. For convenience we will say that a sequence of functions $(f_i)_{i=1}^\infty$ is δ -tapering if $\|2^i \cdot f_i\|_0 \leq \delta$ for all $i \geq 1$.

Lemma 1.2. *Let E be a locally convex subspace of L_0 . For every $\epsilon > 0$ there is a $\delta > 0$ such that if $(f_i)_{i=1}^\infty \subset E$ is δ -tapering then*

$$\mu \left(\bigcup_{i=1}^{\infty} \{x : |f_i(x)| > 1\} \right) \leq \epsilon.$$

Moreover, δ can be chosen to be any positive number such that the closed convex hull of $\{f \in E : \|f\|_0 \leq \delta\}$ is contained in $\{f \in L_0 : \|f\|_0 \leq \frac{\epsilon}{80}\}$.

Proof. Consider the following function on $[0, 1]$:

$$g(t) = \left| \frac{1}{\sqrt{\sum_{j=1}^N a_j^2}} \sum_{k=1}^N a_k r_k(t) \right|,$$

where $a_1, a_2, \dots, a_N \in \mathbb{R}$ and $r_1, r_2, \dots, r_N$ are the first N Rademacher functions. Then from Khinchine's inequality we have

$$\begin{aligned} \int_0^1 g &= \frac{1}{\sqrt{\sum_{j=1}^N a_j^2}} \cdot \left\| \sum_{k=1}^N a_k \cdot r_k \right\|_1 \\ &\geq \frac{1}{\sqrt{\sum_{j=1}^N a_j^2}} \cdot \frac{1}{2} \sqrt{\sum_{k=1}^N a_k^2} \\ &= \frac{1}{2}. \end{aligned}$$

Since the Rademacher functions are orthonormal over $[0, 1]$ we have

$$\begin{aligned} \|g\|_2^2 &= \left\| \sum_{k=1}^N \left(\frac{a_k}{\sqrt{\sum_{j=1}^N a_j^2}} r_k \right) \right\|_2^2 \\ &= \sum_{k=1}^N \left(\frac{a_k}{\sqrt{\sum_{j=1}^N a_j^2}} \right)^2 \\ &= 1. \end{aligned}$$

We now are ready to use Lemma 1.1 with $\alpha = 1/2$ and $\beta = 1/4$:

$$\mu \left(t : \frac{1}{\sqrt{\sum_{j=1}^N a_j^2}} \left| \sum_{k=1}^N a_k r_k(t) \right| \geq \frac{1}{4} \right) \geq \left(\frac{1}{2} - \frac{1}{4} \right)^2 = \frac{1}{16}.$$

Therefore,

$$\mu \left(t : \left| \sum_{k=1}^N a_k r_k(t) \right| \geq \frac{1}{4} \sqrt{\sum_{j=1}^N a_j^2} \right) \geq \frac{1}{16} \quad (*)$$

for $a_1, a_2, \dots, a_N \in \mathbb{R}$.

Let $\epsilon > 0$ be given. Since E is locally convex there is a $\delta > 0$ such that the closed convex hull of $\{f \in E : \|f\|_0 \leq \delta\}$ is contained in $\{f \in L_0 : \|f\|_0 \leq \frac{\epsilon}{80}\}$. Suppose $(f_i)_{i=1}^\infty \subset E$ is δ -tapering. Then for every $N \geq 0$ we have

$$\begin{aligned}
\epsilon/80 &= \int_0^1 \epsilon/80 \, dt \\
&\geq \int_0^1 \left\| \sum_{i=1}^N \frac{1}{2^i} r_i(t) 2^i f_i(x) \right\|_0 \, dt && \text{(local convexity)} \\
&= \int_0^1 \int_0^1 \frac{|\sum_{i=1}^N r_i(t) f_i(x)|}{1 + |\sum_{i=1}^N r_i(t) f_i(x)|} \, dx \, dt \\
&= \int_0^1 \int_0^1 \frac{|\sum_{i=1}^N f_i(x) r_i(t)|}{1 + |\sum_{i=1}^N f_i(x) r_i(t)|} \, dt \, dx && \text{(Tonelli)} \\
&\geq \frac{1}{16} \int_0^1 \frac{\frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}}{1 + \frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}} \, dx. && \text{(by *)}
\end{aligned}$$

So

$$\int_0^1 \frac{\frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}}{1 + \frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}} \, dx \leq 16 \frac{\epsilon}{80} = \frac{\epsilon}{5}.$$

for all $N \geq 1$. Therefore $\mu(x : \sqrt{\sum_{i=1}^N f_i(x)^2} > 1) \leq \epsilon$ for all N . Indeed, suppose not; then

$$\int_0^1 \frac{\frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}}{1 + \frac{1}{4} \sqrt{\sum_{i=1}^N f_i(x)^2}} \, dx > \epsilon \left(\frac{1/4}{1 + 1/4} \right) = \frac{\epsilon}{5}.$$

This is a contradiction. Thus $\mu(x : \sum_{i=1}^N f_i(x)^2 > 1) \leq \epsilon$ for all $N \geq 1$. Letting N go to infinity we get $\mu(x : \sum_{i=1}^\infty f_i(x)^2 > 1) \leq \epsilon$. Finally, since

$$\bigcup_{i=1}^\infty \{x : |f_i(x)| > 1\} \subset \left\{ x : \sum_{i=1}^\infty f_i(x)^2 > 1 \right\}$$

we can conclude that

$$\mu \left(\bigcup_{i=1}^\infty \{x : |f_i(x)| > 1\} \right) \leq \epsilon.$$

□

Lemmas 1.3 and 1.4 find an arbitrarily small set that contains all the ‘unboundedness’ of E .

Lemma 1.3. *Let E be a locally convex subspace of L_0 . Let $\epsilon > 0$ and find $\delta > 0$ so that the closed convex hull of $\{f \in E : \|f\|_0 \leq \delta\}$ is contained in $\{f \in L_0 : \|f\|_0 \leq \frac{\epsilon}{80}\}$. Then if $((f_i^{(k)})_{i=1}^\infty)_{k=1}^\infty \subset E$ is any countable collection of δ -tapering sequences then*

$$\mu \left(\bigcup_{k=1}^\infty \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(k)}(x)| > 1\} \right) \leq \epsilon.$$

Proof. We will start by considering the first n sequences. Let $N_1 < N_2 < \dots < N_{n-1}$. Then

$$(f_i^{(1)})_{i=1}^{N_1} \cup (f_i^{(2)})_{i=N_1+1}^{N_2} \cup \dots \cup (f_i^{(n-1)})_{i=N_{n-2}+1}^{N_{n-1}} \cup (f_i^{(n)})_{i=N_{n-1}+1}^\infty$$

is another δ -tapering sequence. So for all N_{n-1} we have

$$\begin{aligned} \epsilon &\geq \mu \left(\bigcup_{i=1}^{N_1} \{x : |f_i^{(1)}(x)| > 1\} \cup \bigcup_{i=N_1+1}^{N_2} \{x : |f_i^{(2)}(x)| > 1\} \cup \dots \right. \\ &\quad \left. \cup \bigcup_{i=N_{n-2}+1}^{N_{n-1}} \{x : |f_i^{(n-1)}(x)| > 1\} \cup \bigcup_{i=N_{n-1}+1}^\infty \{x : |f_i^{(n)}(x)| > 1\} \right) \\ &\geq \mu \left(\bigcup_{i=1}^{N_1} \{x : |f_i^{(1)}(x)| > 1\} \cup \bigcup_{i=N_1+1}^{N_2} \{x : |f_i^{(2)}(x)| > 1\} \cup \dots \right. \\ &\quad \left. \cup \bigcup_{i=N_{n-2}+1}^{N_{n-1}} \{x : |f_i^{(n-1)}(x)| > 1\} \cup \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(n)}(x)| > 1\} \right). \end{aligned}$$

Let N_{n-1} go to infinity to obtain

$$\begin{aligned} \mu \left(\bigcup_{i=1}^{N_1} \{x : |f_i^{(1)}(x)| > 1\} \cup \bigcup_{i=N_1+1}^{N_2} \{x : |f_i^{(2)}(x)| > 1\} \cup \dots \right. \\ \left. \cup \bigcup_{i=N_{n-2}+1}^\infty \{x : |f_i^{(n-1)}(x)| > 1\} \cup \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(n)}(x)| > 1\} \right) \leq \epsilon. \end{aligned}$$

Repeat this step $n - 2$ times to get

$$\mu \left(\bigcup_{k=1}^n \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(k)}(x)| > 1\} \right) \leq \epsilon.$$

Let n go to infinity to get the desired conclusion,

$$\mu \left(\bigcup_{k=1}^\infty \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(k)}(x)| > 1\} \right) \leq \epsilon. \quad \square$$

In the proof of Lemma 1.4 we use the fact that the space of all Lebesgue measurable subsets of $[0, 1]$ is a complete separable metric space. The distance definition is

$$d(A, B) = \mu(A \Delta B),$$

where $A \Delta B$ stands for the symmetric difference $(A \setminus B) \cup (B \setminus A)$. We consider A and B to be identical if $\mu(A \Delta B) = 0$.

Lemma 1.4. *Let E be a locally convex subspace of L_0 . Let $\epsilon > 0$ and find $\delta > 0$ such that the closed convex hull of $\{f \in E : \|f\|_0 \leq \delta\}$ is contained in $\{f \in L_0 : \|f\|_0 \leq \frac{\epsilon}{80}\}$. Then there is a measurable set A , $\mu(A) \leq \epsilon$, such that if $(f_i)_{i=1}^\infty \subset E$ is any δ -tapering sequence then*

$$\mu \left(\bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i(x)| > 1\} \setminus A \right) = 0.$$

Proof. Let $(f_i^{(t)})_{i=1}^\infty, t \in T$, be the collection of all sequences in E such that $\|2^i \cdot f_i^{(t)}\|_0 \leq \delta$ for all i . T could be an uncountable index set. For each $t \in T$ define

$$A_t = \bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i^{(t)}(x)| > 1\}.$$

$(A_t)_{t \in T}$ is a subspace of the separable metric space consisting of all Lebesgue measurable subsets of $[0, 1]$. So $(A_t)_{t \in T}$ is separable. Let $(A_{t_j})_{j=1}^\infty$ be a countable dense subset. Let

$$A = \bigcup_{j=1}^\infty A_{t_j}.$$

By Lemma 1.3 $\mu(A) \leq \epsilon$. Let $\eta > 0$ and $t \in T$ be given. There is a j such that $\mu(A_{t_j} \Delta A_t) < \eta$ since $(A_{t_j})_{j=1}^\infty$ is dense in $(A_t)_{t \in T}$.

$$\mu(A_t \setminus A) \leq \mu(A_t \setminus A_{t_j}) \leq \mu(A_t \Delta A_{t_j}) < \eta.$$

Since $\eta > 0$ is arbitrary, $\mu(A_t \setminus A) = 0$ for all $t \in T$. □

We are now ready to prove the main theorem. The proof for locally bounded spaces in Kalton, Peck and Roberts [3] was the inspiration for this proof. However, the proofs are quite different in places.

§2. The Lifting Theorem.

Theorem 2.1. *Let E be a locally convex subspace of $L_0[0, 1]$. Let $T : L_0[0, 1] \rightarrow L_0[0, 1]/E$ be a continuous linear operator. Then there is a unique continuous linear operator $S : L_0[0, 1] \rightarrow L_0[0, 1]$ so that $T = QS$, where $Q : L_0[0, 1] \rightarrow L_0[0, 1]/E$ is the quotient map.*

$$\begin{array}{ccc} & & L_0 \\ & \nearrow S & \downarrow Q \\ L_0 & \xrightarrow{T} & L_0/E \end{array}$$

Proof. For each $n = 1, 2, 3, \dots$ find $\delta_n > 0$ so that the closed convex hull of $\{f \in E : \|f\|_0 \leq \delta_n\}$ is contained in $\{f \in L_0 : \|f\|_0 \leq 1/80n\}$, and use Lemma 1.4 to find a measurable set A_n so that

$$(i) \mu(A_n) \leq \frac{1}{n},$$

$$(ii) (f_i)_{i=1}^\infty \subset E \text{ and } \|2^i \cdot f_i\|_0 \leq \delta_n \text{ for all } i \Rightarrow \mu\left(\bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \{x : |f_i(x)| > 1\} \setminus A_n\right) = 0.$$

Without loss of generality we may assume that $\delta_1 \geq \delta_2 \geq \delta_3 \geq \dots$, $\delta_n \rightarrow 0$, and referring to the construction we can take $A_1 \supset A_2 \supset A_3 \supset \dots$. Since T is continuous, for each δ_n we can find $\epsilon_n > 0$ so that $\|f\|_0 \leq \epsilon_n \Rightarrow \|Tf\|_{L_0/E} \leq \delta_n/6$. Without loss of generality we may also assume that $\epsilon_1 \geq \epsilon_2 \geq \epsilon_3 \geq \dots$. For each m and $k = 1, 2, \dots, 2^m$ define

$$\Delta_k^m = \left[\frac{k-1}{2^m}, \frac{k}{2^m} \right).$$

Define

$$\chi_k^m = \chi_{\Delta_k^m} \text{ for } k = 1, \dots, 2^m; m = 1, 2, \dots.$$

Let $v \in L_0$ be given. Define $S(0) = 0$. So we will assume $v \neq 0$. For the next few pages we will work to define $S(v)$. Define $w_k^m = v \cdot \chi_k^m$ for $k = 1, \dots, 2^m$; $m = 1, 2, \dots$. For the time being we will consider m and k to be fixed and look at w_k^m . Let m_0 be the smallest integer so that $1/2^{m_0} \leq \epsilon_1$, and assume $m \geq m_0$. Let $n(m)$ be the largest integer so that $\epsilon_{n(m)} \geq 1/2^m$. Since T is continuous we know that $n(m)$ goes to infinity as m goes to infinity unless T is identically 0. For each $i = 1, 2, \dots$ we can select $g_i \in L_0$ so that $Qg_i = Tw_k^m$ and

$$\|4^i \cdot g_i\|_0 \leq \left(1 + \frac{1}{2^i}\right) \|4^i \cdot Tw_k^m\|_{L_0/E}.$$

If $v = 0$ then $g_i = 0$ for all $i = 1, 2, \dots$. Note that $\|4^i \cdot w_k^m\|_0 \leq 1/2^m \leq \epsilon_{n(m)}$ for all $i = 1, 2, \dots$. So $\|4^i \cdot Tw_k^m\|_0 \leq \delta_{n(m)}/6$ for all $i = 1, 2, \dots$. Therefore $\sigma(Tw_k^m) \leq \delta_{n(m)}/6$.

For $j \geq i \geq 1$,

$$\|4^i(g_i - g_j)\|_0 \leq \|4^i \cdot g_i\|_0 + \|4^j \cdot g_j\|_0 \leq \left(2 + \frac{1}{2^i} + \frac{1}{2^j}\right) \cdot \sigma(Tw_k^m) \leq 3 \cdot \frac{\delta_{n(m)}}{6} < \delta_{n(m)}.$$

Let $f_i = 2^i(g_i - g_{i+1})$. Then $\|2^i \cdot f_i\|_0 \leq \delta_{n(m)}$ for all $i = 1, 2, \dots$. Therefore

$$\mu \left(\bigcap_{l=1}^{\infty} \bigcup_{i=l}^{\infty} \{x : |2^i(g_i - g_{i+1})| > 1\} \setminus A_{n(m)} \right) = 0,$$

that is,

$$\mu \left(\bigcap_{l=1}^{\infty} \bigcup_{i=l}^{\infty} \left\{ x : |g_i - g_{i+1}| > \frac{1}{2^i} \right\} \setminus A_{n(m)} \right) = 0.$$

Let $L(1) = 1$, and for each $p = 2, 3, \dots$ find $L(p) \geq L(p-1)$ such that

$$\mu \left(\bigcup_{i=L(p)}^{\infty} \left\{ x : |g_i - g_{i+1}| > \frac{1}{2^i} \right\} \setminus A_{n(m)} \right) \leq \frac{1}{p}.$$

Define

$$B_p = \bigcup_{i=L(p)}^{\infty} \left\{ x : |g_i - g_{i+1}| > \frac{1}{2^i} \right\} \setminus A_{n(m)}, \quad p = 1, 2, \dots$$

Observe that $B_1 \supset B_2 \supset B_3 \supset \dots$, and $\mu(\bigcap_{p=1}^{\infty} B_p) = 0$. Suppose $x \notin \bigcap_{p=1}^{\infty} B_p$ and $x \notin A_{n(m)}$. We will show that $(g_i(x))_{i=1}^{\infty}$ converges in this case. First there is a p_x such that $x \notin B_{p_x}$. Therefore $|g_i(x) - g_{i+1}(x)| \leq 1/2^i$ for all $i \geq L(p_x)$. Let $\alpha > 0$ be given. Find M such that $2/2^M \leq \alpha$ and $M \geq L(p_x)$. Suppose $j > i \geq M$. Then

$$\begin{aligned} |g_i(x) - g_j(x)| &\leq |g_i(x) - g_{i+1}(x)| + |g_{i+1}(x) - g_{i+2}(x)| + \dots + |g_{j-1}(x) - g_j(x)| \\ &\leq \frac{1}{2^i} + \frac{1}{2^{i+1}} + \dots + \frac{1}{2^{j-1}} \\ &< \frac{2}{2^i} \leq \frac{2}{2^M} \leq \alpha. \end{aligned}$$

So $(g_i(x))_{i=1}^{\infty}$ is a Cauchy sequence in $\mathbb{R}$. For all x define

$$g_k^m(x) = \begin{cases} \lim_{i \rightarrow \infty} g_i(x), & x \notin \bigcap_{p=1}^{\infty} B_p \cup A_{n(m)} \\ 0, & x \in \bigcap_{p=1}^{\infty} B_p \cup A_{n(m)}. \end{cases}$$

g_k^m is the pointwise limit of measurable functions, namely

$$g_i \cdot \chi_{\left(\bigcap_{p=1}^{\infty} B_p\right)^c \cap (A_{n(m)})^c},$$

so g_k^m is measurable. Let $B_k^m = \bigcap_{p=1}^{\infty} B_p$.

We now remember that k and m were arbitrarily chosen, so for each w_k^m we have defined g_k^m and B_k^m for $k = 1, \dots, 2^m$; $m = m_0, m_0 + 1, \dots$. Let

$$B = \bigcup_{k=1}^{\infty} \bigcup_{m=2^k}^{2^{k+1}} B_k^m.$$

$\mu(B) = 0$ since B is the countable union of sets with zero measure. Define

$$S(v) = \lim_{m \rightarrow \infty} \sum_{k=1}^{2^m} g_k^m.$$

It is not immediately clear that $S(v)$ exists. We turn to this question next.

We claim that for almost all $x \notin A_{n(m)}$ ($m \geq m_0$) we have

$$g_p^m(x) = g_{2p-1}^{m+1}(x) + g_{2p}^{m+1}(x).$$

Proof of claim: We know that

$$((g_p^m)_i)_{i=1}^\infty \subset T(w_p^m) \text{ converges in } (A_{n(m)})^c,$$

$$((g_{2p-1}^{m+1})_i)_{i=1}^\infty \subset T(w_{2p-1}^{m+1}) \text{ converges in } (A_{n(m+1)})^c \supset (A_{n(m)})^c,$$

$$((g_{2p}^{m+1})_i)_{i=1}^\infty \subset T(w_{2p}^{m+1}) \text{ converges in } (A_{n(m+1)})^c \supset (A_{n(m)})^c, \text{ and}$$

$$\left((g_{2p-1}^{m+1})_i + (g_{2p}^{m+1})_i \right)_{i=1}^\infty \subset T(w_p^m)$$

because T is additive. (The notation $((g_p^m)_i)_{i=1}^\infty$ simply means the sequence $(g_i)_{i=1}^\infty$ that is associated with w_p^m .) Since $\|4^i (g_p^m)_i\|_0 \leq (1 + 1/2^i) \cdot \|4^i \cdot T(w_p^m)\|_{L_0/E} \leq 2 \cdot \sigma(T(w_p^m)) \leq 2\delta_{n(m)}/6$ we have

$$\|4^i ((g_p^m)_i - (g_{2p-1}^{m+1})_i - (g_{2p}^{m+1})_i)\|_0 \leq \frac{\delta_{n(m)}}{3} + \frac{\delta_{n(m+1)}}{3} + \frac{\delta_{n(m+1)}}{3} \leq \delta_{n(m)}.$$

Therefore

$$\mu \left(\bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \left\{ x : |(g_p^m)_i(x) - (g_{2p-1}^{m+1})_i(x) - (g_{2p}^{m+1})_i(x)| > \frac{1}{2^i} \right\} \setminus A_{n(m)} \right) = 0.$$

So for almost all $x \in (A_{n(m)})^c$

$$\lim_{i \rightarrow \infty} (g_p^m)_i(x) = \lim_{i \rightarrow \infty} ((g_{2p-1}^m)_i(x) + (g_{2p}^m)_i(x)).$$

Thus for almost all $x \in (A_{n(m)})^c$

$$g_p^m(x) = g_{2p-1}^{m+1}(x) + g_{2p}^{m+1}(x),$$

which finishes the proof of the claim.

So the sequence

$$\left(\sum_{k=1}^{2^m} g_k^m \right)_{m=1}^\infty$$

remains essentially fixed in $L_0((A_{n(m)})^c)$ for $r \geq m \geq m_0$. So the sequence converges in $L_0(\bigcup_{m=1}^{\infty} (A_{n(m)})^c) = L_0[0, 1]$, and $S(v)$ is well-defined.

Next we show that $T = QS$. Consider $m \geq m_0$. Then

$$\left\| \left(\sum_{k=1}^{2^m} g_k^m \right) - S(v) \right\|_0 \leq \frac{1}{n(m)},$$

since the two functions are essentially identical except possibly on $A_{n(m)}$ and $\mu(A_{n(m)}) \leq 1/n(m)$. For each k we can find $f_k^m \in T(w_k^m)$ so that

$$\left\| g_k^m|_{(A_{n(m)})^c} - f_k^m|_{(A_{n(m)})^c} \right\|_0 \leq \frac{1}{4^m}.$$

We then have the following inequalities:

$$\left\| \left(\sum_{k=1}^{2^m} g_k^m|_{(A_{n(m)})^c} \right) - \left(\sum_{k=1}^{2^m} f_k^m|_{(A_{n(m)})^c} \right) \right\|_0 \leq 2^m \cdot \frac{1}{4^m} = \frac{1}{2^m},$$

$$\left\| \sum_{k=1}^{2^m} g_k^m - \sum_{k=1}^{2^m} f_k^m \right\|_0 \leq \frac{1}{2^m} + \frac{1}{n(m)},$$

$$\left\| S(v) - \sum_{k=1}^{2^m} f_k^m \right\|_0 \leq \frac{1}{2^m} + \frac{2}{n(m)}.$$

Notice that $1/2^m + 1/n(m) \rightarrow 0$ as $m \rightarrow \infty$. The function

$$\sum_{k=1}^{2^m} f_k^m$$

is an element of $T(v)$. So we can find functions in $T(v)$ that are arbitrarily close to $S(v)$ which means that $S(v) \in T(v)$ since E is closed. That is, $QS(v) = T(v)$.

Next we will show that S is a continuous linear operator. If S is additive and continuous at zero then S must also be homogeneous, and thus linear. So it suffices to show that S is additive and continuous at zero.

S is additive. To see this let $u, v \in L_0$ and let $\alpha > 0$ be given. Find m so that $\mu(A_{n(m)}) \leq \alpha$. (Recall that $\mu(A_{n(m)}) \leq 1/n(m)$.) We will consider $v \cdot \chi_k^m$, $u \cdot \chi_k^m$, and $(u + v) \cdot \chi_k^m$ for an arbitrary k between 1 and 2^m . From our earlier construction we have $(f_i)_{i=1}^{\infty} \subset T(u \cdot \chi_k^m)$ such that $f_i \rightarrow S(u \cdot \chi_k^m)$ on $(A_{n(m)})^c$ and

$$\|4^i \cdot f_i\|_0 \leq \left(1 + \frac{1}{2^i}\right) \|4^i \cdot T(u \cdot \chi_k^m)\|_{L_0(E)},$$

and $(g_i)_{i=1}^\infty \subset T(v \cdot \chi_k^m)$ such that $g_i \rightarrow S(v \cdot \chi_k^m)$ on $(A_{n(m)})^c$ and

$$\|4^i \cdot g_i\|_0 \leq \left(1 + \frac{1}{2^i}\right) \|4^i \cdot T(v \cdot \chi_k^m)\|_{L_0/E},$$

and $(h_i)_{i=1}^\infty \subset T((u+v) \cdot \chi_k^m)$ such that $h_i \rightarrow S((u+v) \cdot \chi_k^m)$ on $(A_{n(m)})^c$ and

$$\|4^i \cdot h_i\|_0 \leq \left(1 + \frac{1}{2^i}\right) \|4^i \cdot T((u+v) \cdot \chi_k^m)\|_{L_0/E}.$$

We have $f_i + g_i \in T((u+v) \cdot \chi_k^m)$ for all $i = 1, 2, \dots$. For $i \geq 1$,

$$\|4^i(f_i + g_i) - 4^i \cdot h_i\|_0 \leq \|4^i \cdot f_i\|_0 + \|4^i \cdot g_i\|_0 + \|4^i \cdot h_i\|_0 \leq (3 + \frac{3}{2^i})\delta_{n(m)}/6 < \delta_{n(m)}.$$

Therefore,

$$\mu\left(\bigcap_{l=1}^\infty \bigcup_{i=l}^\infty \left\{x : |(f_i + g_i) - h_i| > \frac{1}{2^i}\right\} \setminus A_{n(m)}\right) = 0.$$

This implies that $(f_i + g_i)$ and h_i converge to the same function on $(A_{n(m)})^c$. Thus for all $k = 1, \dots, 2^m$, $S(u \cdot \chi_k^m) + S(v \cdot \chi_k^m) = S((u+v) \cdot \chi_k^m)$ on $(A_{n(m)})^c$. Therefore $S(u) + S(v) = S(u+v)$ on $(A_{n(m)})^c$ and $\|S(u) + S(v) - S(u+v)\|_0 \leq \alpha$. Since $\alpha > 0$ was arbitrary we have $S(u) + S(v) = S(u+v)$.

S is continuous at zero. To see this, suppose $(v_j)_{j=1}^\infty$ is a sequence in L_0 such that $v_j \rightarrow 0$. Let $\alpha > 0$ be given. Find m so that $1/n(m) \leq \alpha$. Our set $A_{n(m)}$ then has measure less than α , and $\delta_{n(m)}$ is a positive number such that the closed convex hull of the $\delta_{n(m)}$ -ball in E is contained in the $(\alpha/80)$ -ball in L_0 . There also is an $\epsilon_{n(m)} > 0$ so that $\|f\|_0 \leq \epsilon_{n(m)} \Rightarrow \|Tf\|_{L_0/E} \leq \delta_{n(m)}/6$, and we have $1/2^m \leq \epsilon_{n(m)}$. Let $j \geq 1$ be given. For each $k = 1, \dots, 2^m$ there is a sequence

$$(g_{j,i}^{(k)})_{i=1}^\infty \subset T(v_j \cdot \chi_k^m)$$

such that $g_{j,i}^{(k)} \rightarrow S(v_j \cdot \chi_k^m)$ on $(A_{n(m)})^c$ as $i \rightarrow \infty$ and

$$\|4^i \cdot 4^k \cdot g_{j,i}^{(k)}\|_0 \leq \left(1 + \frac{1}{2^i}\right) \cdot \|4^i \cdot 4^k \cdot T(v_j \cdot \chi_k^m)\|_{L_0/E}.$$

For each $i = 1, 2, \dots$ and $k = 1, \dots, 2^m$ let $f_{i,k} = 2^i \cdot 2^k (g_{j,i}^{(k)} - g_{j,i+1}^{(k)})$. Then $f_{i,k} \in E$ for all i and k and

$$\begin{aligned} \|2^i \cdot 2^k \cdot f_{i,k}\|_0 &= \|4^i \cdot 4^k \cdot (g_{j,i}^{(k)} - g_{j,i+1}^{(k)})\|_0 \\ &\leq \|4^i \cdot 4^k \cdot g_{j,i}^{(k)}\|_0 + \|4^{i+1} \cdot 4^k \cdot g_{j,i+1}^{(k)}\|_0 \\ &\leq 2 \cdot \|4^i \cdot 4^k \cdot T(v_j \cdot \chi_k^m)\|_{L_0/E} \leq \delta_{n(m)} \end{aligned}$$

Using the technique employed in proving Lemma 1.2 we can conclude that

$$\mu \left(\bigcup_{k=1}^{2^m} \bigcup_{i=1}^{\infty} \left\{ x : \left| g_{j,i}^{(k)} - g_{j,i+1}^{(k)} \right| > \frac{1}{2^i} \cdot \frac{1}{2^k} \right\} \right) \leq \alpha.$$

Let the set above be called D (so $\mu(D) \leq \alpha$). Find I such that $2/2^I \leq \alpha$. Then

$$\left\| S(v_j \cdot \chi_k^m) - g_{j,I}^{(k)} \right\|_{L_0((A_{n(m)})^c \cup D^c)} \leq \sum_{i=I}^{\infty} \frac{1}{2^i} \cdot \frac{1}{2^k} = \frac{2}{2^I} \cdot \frac{1}{2^k}.$$

Therefore

$$\left\| S(v_j) - \sum_{k=1}^{2^m} g_{j,I}^{(k)} \right\|_{L_0((A_{n(m)})^c \cup D^c)} \leq \sum_{k=1}^{2^m} \frac{2}{2^I} \cdot \frac{1}{2^k} = \frac{2}{2^I} \leq \alpha,$$

and

$$\left\| S(v_j) - \sum_{k=1}^{2^m} g_{j,I}^{(k)} \right\|_0 \leq 3\alpha.$$

This is true for any $j \geq 1$. Now

$$\left\| \sum_{k=1}^{2^m} g_{j,I}^{(k)} \right\|_0 \leq 2 \sum_{k=1}^{2^m} \left\| 4^I \cdot 4^k \cdot T(v_j \cdot \chi_k^m) \right\|_{L_0/E}.$$

Since T is continuous for each k , $\left\| 4^I \cdot 4^k \cdot T(v_j \cdot \chi_k^m) \right\|_{L_0/E}$ goes to zero as j goes to infinity. Therefore the whole sum goes to zero as j goes to infinity. So $\limsup_{j \rightarrow \infty} \|S(v_j)\|_0 \leq 3\alpha$. However, $\alpha > 0$ was arbitrary, so $\lim_{j \rightarrow \infty} S(v_j) = 0$. That is, S is a continuous linear operator.

Suppose that S' is another continuous linear operator from L_0 to L_0 such that $QS' = T$. Then $Q(S - S') = QS - QS' = T - T = 0$, whence $S - S'$ maps L_0 into the locally convex space E . We conclude that $S = S'$. $\square$

The proof of Theorem 2.1 works with a milder assumption on the subspace E . It does not have to be locally convex - the key assumption is only that given a neighborhood V of 0 there is a smaller neighborhood U so that if $x_n \in U$ then $\sum_{n=1}^N 2^{-n} x_n$ is in V for all N (i.e. E is exponentially galbed in the sense of Turpin [8]). We can generalize further by replacing the sequence (2^{-n}) with a strictly positive term sequence (a_n) such that $\sum a_n < \infty$. By a classical result due to Aoki [1] and Rolewicz [7] we know that locally bounded spaces are locally p -convex for some $p > 0$. Also, if U is locally p -convex then $\sum_{n=1}^N 2^{-(n/p)} U \subset U$ for all N . In this way we can see that the generalized result includes locally bounded subspaces of L_0 .

We can combine Theorem 2.1 with Kwapien's theorem [4].

Theorem 2.2. *Let $S : L_0 \rightarrow L_0$ be a linear operator. Then*

$$S(f)(x) = \sum_{n=1}^{\infty} g_n(x) f(\sigma_n(x))$$

for every $f \in L_0$, where

- (i) each $\sigma_n : [0, 1] \rightarrow [0, 1]$ is a non-singular measurable map,*
- (ii) each g_n is in L_0 ,*
- (iii) for almost all x in $[0, 1]$, $g_n(x) \neq 0$ for only finitely many n .*

Conversely, every map defined in the above way is a linear operator from L_0 to L_0 .

Corollary 2.3. *Let E be a locally convex subspace of L_0 and Q be the quotient map. Then T is an operator from L_0 to L_0/E if and only if $T = QS$ for some S of the form in Theorem 2.2.*

By following the proof of Theorem 4.1 in [2] we have the following corollary.

Corollary 2.4. *Let E and F be subspaces of L_0 , each of which is either locally convex or locally bounded. Then L_0/E is isomorphic to L_0/F if and only if there is an isomorphism S of L_0 to itself such that $S(E) = F$.*

REFERENCES

- [1] T. Aoki, *Locally bounded linear topological spaces*, Proc. Imp. Acad. Tokyo **18** (1942), No. 10.
- [2] N.J. Kalton and N.T. Peck, *Quotients of L_p , $0 \leq p < 1$* , Studia Math. **64** (1979b), 65-75.
- [3] N.J. Kalton, N.T. Peck and J.W. Roberts, *An F -space sampler*, Cambridge University Press, Cambridge, 1984.
- [4] S. Kwapień, *On the form of a linear operator in the space of all measurable functions*, Bull. Acad. Polon. Sci. **21** (1973), 951-954.
- [5] R.E.A.C. Paley and A. Zygmund, *On some series of functions III*, Proc. Camb. Phil. Soc. **28** (1932), 190-205.
- [6] N.T. Peck and T. Starbird, *L_0 is ω -transitive*, Proc. Amer. Math. Soc. **83** (1981), 700-704.
- [7] S. Rolewicz, *On a certain class of linear metric spaces*, Bull. Acad. Polon. Sci. **5** (1957), 471-473.
- [8] P. Turpin, *Dissertationes Mathematicae*, PWN-Polish Scientific Publishers, Warszawa, 1976.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ILLINOIS
URBANA, IL 61801
U.S.A.
E-mail: faber@math.uiuc.edu